

[pra,snowpacs]revtex4 [applemac,inputenc amssymb,amsmath [mathscr]eulscript graphicx

 $k_B[1]\langle 1 \rangle LH$

document

arXiv:quant-ph/0211072

A simple quantum heat engine

Jacques Arnaud

arnaudj2@wanadoo.fr

Mas Liron, F30440 Saint Martial, France

Laurent Chusseau

chusseau@univ-montp2.fr

<http://www.opto.univ-montp2.fr/~chusseau>Centre d'lectronique et de Micro-optoelectronique de Montpellier, UMR 5507 CNRS, Universit Montpel-
lier II, F34095 Montpellier, France

Fabrice Philippe

[Also at]MIAp, Universit Paul Valry, F34199 Montpellier, France

Fabrice.Philippe@univ-montp3.fr

Laboratoire d'Informatique de Robotique et de Microelectronique de Montpellier, UMR 5506 CNRS, 161
Rue Ada, F34392 Montpellier, France

abstract Quantum heat engines employ as working agents multi-level systems instead of gas-filled cylinders. We consider particularly two-level agents such as electrons immersed in a magnetic field. Work is produced in that case when the electrons are being carried from a high-magnetic-field region into a low-magnetic-field region. In watermills, work is produced instead when some amount of fluid drops from a high-altitude reservoir to a low-altitude reservoir. We show that this purely mechanical engine may in fact be considered as a two-level quantum heat engine, provided the fluid is viewed as consisting of n molecules of weight one and $N - n$ molecules of weight zero. Weight-one molecules are analogous to electrons in their higher energy state, while weight-zero molecules are analogous to electrons in their lower energy state. More generally, fluids consist of non-interacting molecules of various weights. It is shown that, not only the average value of the work produced per cycle, but also its fluctuations, are the same for mechanical engines and quantum (Otto) heat engines. The reversible Carnot cycles are approached through the consideration of multiple sub-reservoirs.